

THE IMPACT OF A CLOSE STELLAR ENCOUNTER ON THE EDGEWORTH-KUIPER BELT

ALICE C. QUILLEN

Department of Physics and Astronomy, University of Rochester, Rochester, NY 14627
 aquillen@pas.rochester.edu

DAVID E. TRILLING, &

University of Pennsylvania, 209 South 33rd Street, Philadelphia, PA 19104
 trilling@hep.upenn.edu

ERIC G. BLACKMAN

Department of Physics and Astronomy, University of Rochester, Rochester, NY 14627
 blackman@pas.rochester.edu
Draft version February 2, 2008

ABSTRACT

We numerically investigate the possibility that a close stellar encounter could account for the high inclinations of the Kuiper belt, as originally proposed by Ida, Larwood and Burkert, however we consider encounters with pericenters within those explored by previous works. A star of mass $0.2M_{\odot}$ entering the solar system on a nearly parabolic, low inclination, retrograde orbit with perihelion of 50 AU can scatter $\sim 30\%$ of the Kuiper belt into moderate inclination ($i > 10^{\circ}$) and eccentricity orbits, while leaving the rest of the belt at low eccentricities and inclinations. This scenario provides a possible explanation for the dichotomy of the Kuiper belt, accounting for the scattered and classical populations, including objects with high eccentricities and inclinations. High eccentricity objects that were not placed into resonance with Neptune are more likely to be removed from the system by subsequent long timescale evolution. Consequently this scenario suggests that some Plutinos could have originally been objects in circular orbits which were scattered into resonant high eccentricity and high inclination orbits by the stellar encounter. Objects could have been placed into stable resonant regions with Neptune by the encounter. However long lived regions of dynamical stability should differ from those selected by resonance capture. This scenario predicts the presence of objects in resonances with Neptune that are unlikely to have been captured via Neptune's migration.

Subject headings: Kuiper Belt — planetary systems: protoplanetary disks

1. INTRODUCTION

The Edgeworth Kuiper Belt (KB) is often discussed in terms of three dynamical families; 1) the classical Kuiper Belt objects which occupy semi-major axes $41 \lesssim a \lesssim 46$ AU and which have low eccentricities $e < 0.25$, 2) the resonant Kuiper Belt occupying the 3:2 and 2:1 mean motion resonances with Neptune, denoted the Plutinos and Twotinos, respectively, and 3) the scattered Kuiper belt occupying extreme orbits with high eccentricities (e up to 0.6) (e.g. Trujillo & Brown 2002). Most of the mass in the Kuiper belt appears to reside in the classical Kuiper belt (Luu & Jewitt 2002). All three dynamical families contain objects with surprisingly high inclinations ($i > 15^{\circ}$; Trujillo & Brown 2002; Luu & Jewitt 2002; Brown 2001). Current compilations suggest a bimodal distribution, with $\sim 20\%$ of the Kuiper belt objects in low-inclination orbits of a few degrees and the remainder in highly inclined orbits with $i \gtrsim 20^{\circ}$ (Brown 2001). Scenarios for the formation and evolution of the Kuiper Belt should explain the different Kuiper Belt Object (KBO) populations.

Several mechanisms have been proposed to account for the high KBO inclinations, including scattering by Neptune during Neptune's migration (Gomes 2003; Levison & Morbidelli 2003), passage through vertical resonances during Neptune's migration or during the depletion of the solar nebula (Nagasawa & Ida 2000; Malhotra 1998), wave excitation (Ward 2003), perturbations from passing stars

(Ida et al. 2000; Kobayashi & Ida 2001), and scattering by planets and planetesimals (Thommes et al. 2002; Petit et al. 1999; Kenyon 2002; Brunini & Melita 2002). While long term evolution can remove KBOs with low inclinations, it cannot explain the existence of classical KBOs with extremely high inclinations of 30° (Kuchner et al. 2002). Current surveys appear to rule out the possibility of a distant planetary embryo (Morbidelli et al. 2002).

In this paper we focus on the possibility that perturbations from a stellar encounter could account for the high KBO inclinations, as proposed by Ida et al. (2000). The probability of a stellar encounter that passes with a few hundred AU of the Sun at its current position is extremely unlikely. However, most stars are formed in stellar groups or clusters (Lada & Lada 2003), associated for $10^7 - 10^9$ years until they are tidally disrupted by molecular clouds and structure in the Milky Way disk (Lada & Lada 2003; Terlevich 1987; Theuns 1992; Bergond et al. 2001). Young stellar groups and clusters, can have stellar densities up to 10^5 times higher than the Solar neighborhood (Lada & Lada 2003; Hillenbrand & Hartmann 1998). The Sun could have experienced a close stellar encounter when it was still in its birth cluster, (Ida et al. 2000; Adams & Laughlin 2001; Gaidos 1995). Other disk systems such as Beta Pictoris and HD 100546 probably have suffered close stellar encounters (Larwood & Kalas 2001; Quillen & Thorndike 2002). Alternatively, the Sun could have been

part of a wide binary which was later disrupted (e.g., as Furlan et al. 2003).

As shown by Larwood & Kalas (2001) in their study of the Beta Pictoris system, and others (e.g. Kobayashi & Ida 2001) a close stellar encounter can scatter planetesimals. For an equal mass parabolic encounter with pericenter distance q , planetesimals with semimajor axis $a \gtrsim q/2$ are scattered, whereas those with $a \lesssim q/2$ are uniformly perturbed into higher eccentricity and inclination orbits (Larwood & Kalas 2001; Kobayashi & Ida 2001). This division depends on the timescale of the encounter compared to the orbital rotation period of the planetesimals. In the inner disk, the planetesimals rotate quickly compared to the encounter time and the collision can be treated with a secular approximation, averaging over the entire orbit, as done in the appendix by Kobayashi & Ida (2001). However, objects with larger semimajor axes are moving at speeds similar to the encountering star (assuming a parabolic orbit). Consequently the resulting perturbation on outer planetesimals is strongly dependent on their longitudes, as shown explicitly by Larwood & Kalas (2001). If the collision is very fast compared to the rotational timescale of the planetesimals, as would be the case if the star was on a highly hyperbolic orbit, then the impulse approximation can be used to estimate the perturbations caused by the encounter (e.g., Eggers & Woolfson 1996; Pfalzner 2003).

Previous work considering the effect of a stellar encounter on our solar system has considered parabolic orbits with pericenter of size 100-200 AU (Ida et al. 2000; Kobayashi & Ida 2001). At this distance the velocity of the perturber at pericenter is similar to the rotational velocity of KBOs. For such an encounter the Kuiper belt is in a transition region. The KBOs experience increases in mean eccentricity and inclination which are dependent on their initial semimajor axes. Because the final KBO orbits in this region also depend on the initial longitude, the eccentricity and inclination distribution broadens following the encounter. Ida et al. (2000); Kobayashi & Ida (2001) restricted their exploration to fairly large stellar pericenters $q > 100$ AU, which minimized the eccentricity variations in the KBOs caused by the encounter. Ida et al. (2000); Kobayashi & Ida (2001) suggested that Neptune's migration followed the encounter. KBOs with initial eccentricities above 0.15 are unlikely to be captured in the 3:2 mean motion resonance with a migrating Neptune (Malhotra 1995; Hahn & Malhotra 1999). The encounters considered by Ida et al. (2000); Kobayashi & Ida (2001) were consistent with a scenario that included migration of Neptune since the encounters did not induce high KBO eccentricities.

In this paper we extend the previous explorations by Ida et al. (2000); Kobayashi & Ida (2001) to include stars on orbits with smaller pericenters. In this regime the planetesimal perturbations are strongly dependent on the planetesimal longitude during the encounter. Consequently we expect a larger spread in the resulting planetesimal planetesimal orbital properties, as seen in the outer parts of the planetesimal disks simulated by Larwood & Kalas (2001); Ida et al. (2000); Kobayashi & Ida (2001). Here we explore a regime where the Kuiper Belt is more effectively scattered by the encounter than previously works

have considered.

2. CONSTRAINTS ON PARAMETER SPACE OF THE ENCOUNTER

In this section we discuss preliminary constraints on the orbit, mass and initial velocity of the stellar impactor. We maintain our hypothesis that the Kuiper Belt is strongly scattered by the encounter, but that it must retain low eccentricity and inclination objects as are found in the classical Kuiper Belt. Furthermore we assert that Uranus's orbit is not strongly perturbed by the encounter.

The Sun/stellar encounter is described by 5 free parameters, the mass of the star compared to the Sun, $M_*/M_\odot$, the velocity of the star distant from the Sun, v_∞ , the distance of closest approach from the Sun or perihelion of the star, q , the inclination of the star's orbit with respect to the ecliptic, i_* , and the argument of perihelion, ω_* . For a pictorial view of these angles see Fig. 1 which is based on Fig. 1 by Kobayashi & Ida (2001). The velocity of the incoming star is related to the eccentricity of the orbit e_* , where $v_\infty^2 = (e_* - 1) \left(\frac{G(M_\odot + M_*)}{q} \right)$ and $e_* > 1$ for a hyperbolic orbit.

For a star to scatter a planetesimal effectively, the star's velocity must be greater than the circular velocity of the planetesimal. The velocity of a star in a hyperbolic orbit at perihelion is

$$v_q^2 = (1 + e_*) \frac{G(M_\odot + M_*)}{q}. \quad (1)$$

We compare the star's velocity at perihelion with the velocity, v_c , of a particle in a circular orbit with semi-major axis a ; $v_c = \sqrt{GM_\odot/a}$. Allowing Neptune to be in the scattered region, we require $v_c \gtrsim v_q$ at Neptune's $a = 30$ AU which implies that

$$q/\sqrt{1 + e_*} \lesssim 30 \text{ AU}. \quad (2)$$

This sets pericenter $q \lesssim 50 \text{ AU}$ if the star is on a parabolic orbit and if Neptune is in the scattered region.

We now constrain e_* based on the most probably stellar incoming velocity. Collisions are unlikely from field stars, so we expect an incoming stellar velocity of order a few km/s, typical of the velocity dispersion in young star clusters which have not yet dispersed or disrupted.

$$v_\infty^2 = (e_* - 1) \left(\frac{G(M_\odot + M_*)}{q} \right) \quad (3)$$

The circular velocity is 4.2 km/s at a semi-major axis of 50 AU, of order that which might be found in a young unbound cluster. Since we don't expect v_∞ to be greater than a few km/s if the perturber is in a birth cluster, we infer that the encounter was likely to be nearly parabolic, ($e_* \lesssim 2$). A hyperbolic orbit with a small pericenter from a faster field star would be possible, but it is much less likely than a parabolic one from a star in a dense birth cluster. If the encounter was a result of an interaction with a binary, then the velocity of the encounter would depend on the nature of the binary and how it was disrupted.

We can set a condition on the orbit of the impactor by specifying that Uranus does not significantly increase in inclination or eccentricity following the encounter. Uranus is at small enough semimajor axis that we may assume that it moves quickly compared to the timescale of the encounter. To estimate the affect of the encounter on Uranus

we use equations 15 and 16 from Kobayashi & Ida (2001) which are appropriate for parabolic encounters in a secular approximation.

$$i \approx \frac{3\pi}{8\sqrt{2}} \frac{M_*/M_\odot}{\sqrt{1+M_*/M_\odot}} \left(\frac{a}{q}\right)^{3/2} |\sin 2i_*| \quad (4)$$

Based on Uranus's inclination, 0.77° , and semimajor axis, 19.2 AU, and assuming an encounter with $q \sim 50$ AU we find

$$M_* \sin 2i_* \lesssim 0.087. \quad (5)$$

If the star was low mass $M_* \sim 0.1$, then a reasonably large range of inclinations is allowed. Since Uranus's eccentricity increase is $\propto (a/q)^{5/2}$ (Kobayashi & Ida 2001), Uranus's low eccentricity is less of a constraint than its low inclination. If Uranus was not strongly perturbed by the encounter, then the stellar impactor was likely to be a low mass star or on a low inclination orbit.

For particles with semi-major axes similar to the pericenter distance of the encounter, the perturbation caused by the encounter is strongly dependent upon the particle's longitude. Since these particles move slower than the star, we can use the impulse approximation to estimate the size of the perturbations. Each particle undergoes a change in its velocity of order

$$|\Delta V| \sim \frac{2GM_*}{bv_q} \quad (6)$$

where b is the distance between the object and the star's closest approach to it, and v_q is the velocity of the star at pericenter (e.g., Binney & Tremaine 1987; Eggers & Woollson 1996). The velocity change is in the direction toward the star at its position of closest approach. Using cylindrical coordinates (R, z) where R is in the ecliptic, the position of the star's pericenter is $R = q\sqrt{\cos^2 \omega_* + \sin^2 \omega_* \cos^2 i_*}$, $z = q \sin \omega_* \sin i_*$. For an object opposite pericenter (180° away)

$$\frac{\Delta v_z}{v} \sim \sin i \sim \left(\frac{M_*}{M_\odot}\right)^{1/2} \frac{\sqrt{2aq}}{a+q} \sin \omega_* \sin i_* \quad (7)$$

To allow the inclination of particles opposite pericenter to remain low, we restrict the encounter to low arguments of perihelion and low inclinations.

3. SIMULATIONS OF HYPERBOLIC ENCOUNTERS

From the previous section we found that if Neptune is in the scattered region, then $q \lesssim 50$ AU. If the incoming star is part of the Sun's birth cluster then we expect $e_* \lesssim 2$. To maintain Uranus's low eccentricity and inclination, we require low inclination orbits and low mass stars. To ensure that the Kuiper Belt retains a low eccentricity and inclination population we require orbits with low inclinations and low arguments of perihelion. We take these estimates as guidelines for our numerical integrations.

To investigate the actual affect of a such a close stellar encounter on a primordial cold Kuiper belt, we numerically integrated the two body Sun/stellar gravitational system. Our code is a conventional Burlisch-Stoer numerical scheme which considers only the force from Newtonian gravity. Simultaneously during the encounter we integrate 1000 low mass particles (planetesimals) per run which were

initially placed in circular orbits in the ecliptic. These particles are influenced by the gravitational force from the Sun and star but do not themselves act on the Sun or star. The integrations were begun with the star at 500 AU from the Sun, and ended when the star reached an equivalent distance from the Sun following the encounter. We began our simulations with planetesimals distributed randomly in semi-major axis between 30 and 70 AU and randomly chosen mean longitudes. Following the encounter, the final semi-major axes, eccentricities and inclinations of the particles were tabulated.

Based on our constraints from the previous section, we explored planetesimal eccentricity and inclination distributions for encounters spanning $0.1 < M_*/M_\odot < 0.5$, $40 < q < 70$ AU with differing orbital inclinations and arguments of perihelion (i_*, ω_*) . Since we expect that the encounter could have been in the Sun's birth cluster, we restrict the incoming stellar velocity v_∞ to less than a few km/s, equivalent to nearly parabolic encounters, $e_* < 2$. In this region of parameter space, we searched for final planetesimal distributions which contained both scattered and classical type KBO population analogs.

In Fig. 2 we show the planetesimal distribution following a retrograde encounter with a star of $M_*/M_\odot = 0.2$, $e_* = 1.4$, pericenter $q = 40$ AU, inclination $i_* = 170^\circ$ and argument of perihelion $\omega_* = 20^\circ$. In Fig. 2, histograms are also shown for all particles with final semi-major axes between 33 and 50 AU. Examination of this figure shows that the inclination and eccentricity distribution following the encounter is wider at larger semi-major axes as expected from a comparison between the timescale of the encounter with the orbital period of the objects. We see that a significant fraction, but not all, of the belt is scattered, indicating that the final eccentricity and inclination of an object depends on its longitude during the collision (as shown explicitly by Larwood & Kalas 2001). About 30% of objects with final semimajor axes between 33 and 50 AU have inclinations above 10° and eccentricities above 0.1. However, low eccentricity ($e < 0.05$) and low inclination ($i < 6^\circ$) objects remain in the planetesimal distribution.

Fig. 2 can be compared to the distribution of known Trans-Neptunian objects (as listed by the Minor Planet Center Jan. 7, 2004¹) which is given in the same format in Fig. 3. To ensure accurate orbits, only objects that have been observed over more than one opposition have been plotted. We must keep in mind that the known population of KBOs is likely to be biased by the systematics of the observations (e.g., as discussed by Bernstein et al. 2004; Brown 2001). The KBO has also had ~ 4 billion years to evolve dynamically following the hypothetical collision considered here (e.g., as explored by Kuchner et al. 2002).

While the simulation of the retrograde encounter shown in Fig. 2 does not exactly match the known KBO population (the most obvious mismatch being the radial distribution), it successfully predicts the combined presence of objects with high inclinations and eccentricities while retaining a population with low inclinations and eccentricities, as required by the known KBO distribution. The inclination and eccentricity distribution following the en-

¹ <http://cfa-www.harvard.edu/iau/lists/TNOs.html>

counter have high inclination and eccentricity tails, similar to those seen in the KBO comparison histograms. An object at the location of Neptune (30AU) can remain in a low eccentricity and inclination orbit, so Neptune itself could have survived the encounter with minimal changes in its orbital elements.

The distribution of orbital elements for the retrograde encounter can be contrasted, for example, with the prograde encounter shown in Fig. 4 which does not exhibit as broad a distribution of eccentricities and inclinations. The simulation (Fig. 2) we chose to directly compare to the current KBO population (Fig. 3) corresponded to a low inclination retrograde encounter. A prograde encounter can more effectively perturb the disk because the collision effectively lasts longer in the frame rotating with each planetesimal. We also find prograde collisions which are capable of scattering the disk, and an example of one is shown in Fig. 4. While low inclination retrograde encounters fail to produce very low inclination objects in the belt, we find that the low inclination prograde close encounters fail to produce low eccentricity objects, requiring subsequent evolution to account for them, including the low eccentricity of Neptune itself. In addition, we have had difficulty identifying a region of parameter space in which a low eccentricity and inclination population is retained but a significant fraction of the belt is scattered.

We consider what effects subsequent evolution could have on the resulting planetesimal distribution shown in Fig. 2 for the retrograde encounter. Scattered objects at high eccentricities that were not placed in resonances with Neptune would be quickly removed from the system because they cross Neptune's orbit and so could be scattered by Neptune. High eccentricity objects at the location of Pluto (40AU) would be removed, leaving a population of Plutinos which include high eccentricity and inclination objects. Alternatively because there remains a population of low eccentricity objects, subsequent migration by Neptune can resonantly capture objects into the 3:2 resonance accounting for the Plutinos. Because the low eccentricity objects include objects at high inclinations, the resulting captured Plutinos would also include high inclination objects.

The simulation shown in Fig. 2 does not necessarily require Neptune to migrate to account for the Plutinos since non resonant high eccentricity objects at 40 AU will be removed by subsequent scattering by Neptune. If we consider models which include Neptune's migration, then a closer (smaller pericenter) encounter than simulated in Fig. 2 would be required. This follows because low eccentricity objects captured at smaller semi-major axes must include high inclination objects such as are found past 40 AU in this simulation.

While no objects are found at extremely low inclinations in the retrograde simulation, (Fig. 2), subsequent evolution may broaden the inclination distribution, reducing the inclinations of some objects. We began the simulations with zero inclination and eccentricity objects. Had we started the simulation with a distribution of inclinations and eccentricities, the final distribution would have included lower inclination objects. If the stellar orbit has a higher inclination than the simulation shown in 2, then following the encounter the mean inclination is higher, how-

ever more objects remain at low eccentricities, and the eccentricity distribution is narrower.

Deep observational searches suggest that the Kuiper Belt is truncated at 50 AU (Allen et al. 2001) and does not contain a large population of "cold" or near zero eccentricity objects (Allen et al. 2002). Our simulated encounters do not truncate the disk, but instead increase the radial distribution of any previously defined edge (e.g. as shown by (Larwood & Kalas 2001) in similar simulations for the Beta Pic disk). It is possible that an edge in the KBO distribution existed prior to the stellar encounter. To investigate the effect of this edge, we ran encounter simulations with a predefined edge at 45 AU in the planetesimal distribution prior to the encounter. Following the encounter, the planetesimal distribution is shown in Fig. 5. The simulated encounter was retrograde with $M_* = 0.25M_\odot$, $e_* = 1.3$, $i_* = 170^\circ$, $\omega_* = 20^\circ$, and $q_0 = 40\text{AU}$. The resulting planetesimal distribution exhibits a rise in both mean eccentricity and inclinations with semi-major axis for objects past 45 AU. A rise in mean eccentricity is exhibited by the known KBOs shown in Fig. 3, however the known KBO distribution lacks an increase in mean inclination in the region same region. Consequently the observed increase in mean eccentricity with semimajor axis seen in the KBO population is unlikely to be explained by an encounter perturbing a pre-existing disk edge. Since the brightness of an object depends on its distance from the Sun to the fourth power, high eccentricity KBO's are more likely to be discovered when they are near perihelion, even taking into account the time they spend at small distances from the Sun compared to that spent at large distances. This selection effect may have reduced the number of known low eccentricity KBOs objects at large semimajor axis.

4. DISCUSSION AND SUMMARY

In this paper we have numerically investigated the effect of a close stellar encounter on the primitive Kuiper Belt. We have extended our study to collisions with lower mass stars and smaller pericenters than have the pioneering studies of Ida et al. (2000); Kobayashi & Ida (2001). We find that low mass, low inclination encounters can produce a dichotomy in the resulting planetesimal distribution, containing both scattered objects with high eccentricities or high inclinations as well as low inclination and low eccentricity objects, similar to what is seen in the Kuiper Belt. In particular a $0.2M_\odot$ star passing with a pericenter of 45 AU from the Sun, in a retrograde, highly inclined orbit produces inclination and eccentricity distributions with high end tails, similar to those seen in the KBO distributions.

To more thoroughly test such a scenario it is necessary to numerically evolve the scattered population for the lifetime of the solar system as done for the classical KBO population by Kuchner et al. (2002). High eccentricity objects at 40AU which were not placed in resonance with Neptune are likely to be removed from the Kuiper Belt by scattering with Neptune. Consequently the 3:2 resonance represents a region of stability; objects placed in this resonance could stay there for the lifetime of the solar system. Objects placed in this resonance by the encounter could explain the high inclination Plutino population. Ob-

jects currently residing in resonances that have low capture probabilities such as the 5:2 resonance (Chiang et al. 2003) might be more naturally explained with a scattering scenario rather than a Neptune migration scenario. Alternately one could regard the existence of such objects as a consequence or prediction of the impact scenario explored here. Since long lived regions of dynamical stability differ from those selected by resonance capture, study of future measurements of the of KBO orbital distribution may support or reject a scenario which includes a close stellar encounter.

Low eccentricity objects remain in the belt following the encounter, which implies that there remains a population which can be captured into resonance by subsequent migration by Neptune (e.g., Nagasawa & Ida 2000), however Neptune’s migration may not be needed to explain the highest Plutino eccentricities. This may reduce the extent of migration required by recent migration models (Gomes 2003; Levison & Morbidelli 2003). Hahn & Malhotra (2003) have suggested that the resonant KBO populations might be better explained by Neptune’s migration into a previously “hot” Kuiper Belt, a situation which could occur after the encounter we have simulated here.

Following the stellar encounter, we expect many objects to be removed from the belt by dynamical evolution, suggesting that there was a period of time when short period comets were produced in great numbers. One interesting possibility is that this epoch corresponded to the period of “Late Heavy Bombardment” when the solar system was a few $\times 10^8$ years old. This possibility would require a cometary origin for the lunar impacts associated with the Late Heavy Bombardment Era (for further discussion on the cometary vs. asteroidal origin see Levison et al. 2000, 2001). Because of the large mass of material likely to be sent into the inner solar system by Neptune, this epoch is also very likely to be the time during which Neptune migrated. Because of the narrowness of Neptune’s mean motion resonances, a close stellar encounter probably did not occur after the migration of Neptune. This follows because such an encounter would have effectively removed objects from resonance. The encounter explored here would also place objects in the 2:1 resonance with Neptune, and so may not naturally account for the low fraction of Twotinos compared to Plutinos unless there was a pre-existing edge to the Kuiper Belt.

While a close encounter with a low mass star could account for the inclination distribution and dichotomy of orbital properties seen in the Kuiper Belt, as seen in our simulations, it does not truncate the disk. Because they pull tidal arms outward, stellar encounters do not tend to sharply truncate disks (e.g. Pfalzer 2003). If the Kuiper Belt contains an edge at 50 AU as suggested by (Allen et al. 2001), then it is much more likely to have been caused by a companion, either stellar or planetary. We note that the encounters simulated here did remove $\sim 10\%$ of the planetesimals in the region 30–70AU objects from the Solar system. The encounter could have ejected a primordial planet at a semimajor axis of ~ 50 AU, leaving only its signature behind in the KBO distribution.

If a stellar encounter did perturb the young Kuiper Belt, then the limits on the likely stellar birth cluster of

the Sun can be somewhat relaxed from the lower limits placed by Adams & Laughlin (2001), as suggested by (de la Fuente Marcos & de la Fuente Marcos 2001). In particular a higher stellar density birth cluster, similar to that in the Orion nebula with a few times 10^4 stars pc^{-3} (Hillendbrand & Hartmann 1998), or a longer lasting open cluster, would be required to make the hypothetical collision explored here probable during the lifetime of the cluster. The probability P , of an object of cross section σ undergoing a collision while moving through a stellar population of number density n , with velocity dispersion $v \sim 1$ km/s during time t is $P = n v \sigma t$ (e.g., Adams & Laughlin 2001). Due to gravitational focusing, the impact parameter b , of an encounter is larger than the pericenter distance q . For an encounter with a small pericenter (when $\sqrt{G(M_\odot + M_*)}/q > v_\infty$), the impact parameter is $\frac{b}{q} \sim \frac{\sqrt{G(M_\odot + M_*)/q}}{v_\infty}$. For an encounter with the Sun with pericenter $q \sim 50$ AU, and $v_\infty = 1$ km/s, we estimate a cross section $\sigma = \pi b^2 \sim 3 \times 10^4 \text{AU}^2$. In the Sun’s birth cluster the probability that such an encounter occurred during time t

$$P \sim 0.16 \left(\frac{n}{2 \times 10^4 \text{pc}^{-3}} \right) \left(\frac{v}{1 \text{km s}^{-1}} \right) \left(\frac{\sigma}{3 \times 10^4 \text{AU}^2} \right) \left(\frac{t}{110 \text{Myr}} \right). \quad (8)$$

We therefore find that a collision proposed here would have been only a moderately improbable event (15 percent level) if the Sun were born in a dense cluster similar to that in the Orion nebula. Here we have only considered the probability of a close collision and have not restricted the orientation of the encounter with respect to the ecliptic. Had we taken into account the narrow region of allowed orbital inclinations and arguments of perihelion, our proposed encounter would be about 100 times less probable. The formation time of the KBOs ($\gtrsim 10^8$ years) is longer than that used in the above estimate, (Kenyon & Luu 1998; Stern & Colwell 1997), suggesting that the Sun resided in a longer lived and more massive birth cluster, such as an open cluster. A longer timescale would allow the close encounter to occur after the formation of the KBOs. Alternatively the Sun could have been part of a wide binary, and the close encounter could have occurred when the binary was disrupted (e.g., one of the components of T Tauri has just recently been disrupted, see Furlan et al. 2003).

Currently there are number of examples of young systems which show clear evidence of external perturbers, including the AeBe stars HD 141569A which has spiral structure and asymmetries excited by its binary companion HD 141569B,C (Augereau & Papaloizou 2003; Quillen et al. 2004), and HD 100546 which exhibits spiral structure which could have been excited by star at a pericenter of about 500 AU (Quillen et al. 2004; Grady et al. 2001). The tilt in Bet Pictoris’s disk has been explained with an stellar encounter with a similar sized pericenter (Larwood & Kalas 2001). Structure seen in submillimeter emission of older systems have recently been interpreted in terms of high eccentricity outer planets. For example models for the submillimeter distribution for Vega include an outer high eccentricity Jupiter mass planet (Wilner et al. 2002) and that for Epsilon Eridani include a Neptune mass moderate eccentricity planet (Quillen & Thorndike 2002). While

the hypothetical encounter with our solar system explored here did not significantly increase Neptune's eccentricity, it easily could have, had the incoming star approached the Sun from a different orientation. This suggests a possible scenario which could account for the observed morphology of the the dusty debris disks which require outer eccentric planets. Future study may determine where and when close stellar encounters are likely to affect young solar systems as well as test and further explore the possibility that such an encounter occurred in our own Solar system.

We thank S. Thorndike, E. Chiang, E. Ford, P. Varniere, and A. Frank for helpful discussions. This material is based on upon work supported by the National Aeronautics and Space Administration under Grant No.xxx issued through the Origins of Solar Systems program. This research was supported in part by the National Science Foundation to the KITP under Grant No. PHY99-07949.

REFERENCES

- Adams, F. C., Laughlin, G. 2001, *Icarus*, 150, 151
- Allen, R. L., Bernstein, G. M., & Malhotra, R. 2001, *ApJ*, 549, L241
- Allen, R. L., Bernstein, G. M., & Malhotra, R. 2001, *AJ*, 124, 2949
- Augereau, J. C., & Papaloizou, J. C. B. 2003, *A&A*, in press, (astro-ph/0310732)
- Bergond, G., Leon, S., & Guibert, J. 2001, *A&A*, 377, 462
- Bernstein, G. M., Trilling, D. E., Allen, R. L. Brown, M. E., Holman, M., & Malhotra, R. 2004, *AJ*, in press
- Binney, J., & Tremaine, S. 1987, *Galactic Dynamics*, Princeton University Press, Princeton, NJ
- Brown, M. E. 2001, *AJ*, 121, 2804
- Brunini, A., & Melita, M. D. 2002, *Icarus*, 160, 32
- Chiang, E. I., Jordan, A. B., Millis, R. L., Buie, M. W., Wasserman, L. H., Elliot, J. L., Kern, S. D., Trilling, D. E., Meech, K. J., & Wagner, R. M. 2003, *AJ*, 126, 430
- Eggers, S., Woolfson, M. M. 1996, *MNRAS*, 282, 13
- de la Fuente Marcos, C., & de la Fuente Marcos, R. 2001, *A&A*, 371, 1097
- Furlan, E., Forrest, W. J., Watson, D. M., Uchida, K. I., Brandl, B. R., Keller, L. D., & Herter, T. L. 2003, *ApJ*, 596, L87
- Gaidos, E. J. 1995, *Icarus*, 114, 258
- Gomes, R. S. 2003, *Icarus*, 161, 404
- Grady, C., et al. 2001, *AJ*, 122, 3396
- Hahn, J. M., & Malhotra, R. 1999, *AJ*, 117, 3041
- Hahn, J. M., & Malhotra, R. 2003, American Astronomical Society, Division for Planetary Sciences Meeting, #35, #39.08
- Hillenbrand, L. A., & Hartmann, L. W. 1998, *ApJ*, 492, 540
- Ida, S., Larwood, J., & Burkert, A. 2000, *AJ*, 528, 351
- Kenyon, S. J. 2002, *PASP*, 114, 265
- Kenyon, S. J., & Luu, J. X. 1998, *AJ*, 115, 2136
- Kobayashi, H., & Ida, S. 2001, *Icarus*, 153, 416
- Kuchner, M. J., Brown, M. E., & Holman, M. 2002, *AJ*, 123, 1221
- Lada, C. J., & Lada, E. A. 2003, *ARA&A*, 41, 57
- Larwood, J. D. & Kalas, P. G. 2001, *MNRAS*, 323, 402
- Levison, H. F. & Morbidelli, A. 2003, *Nature*, 426, 419
- Levison, H. F., Duncan, M. J., Zahnle, K., Holman, M., & Dones, L. 2000, *Icarus*, 143, 415
- Levison, H. F., Dones, L., Chapman, C. R., Stern, S. A., Duncan, M. J., & Zahnle, K. 2001, *Icarus*, 151, 286
- Luu, J. X., & Jewitt, D. C. 2002, *ARA&A*, 40, 63
- Malhotra, R. 1995, *AJ*, 110, 420
- Malhotra, R. 1998, *Lunar Planet. Sci. Conf.*, 29, 1476
- Morbidelli, A., Jacob, C., & Petit, J.-M. 2002, *Icarus*, 157, 241
- Nagasawa, M., & Ida, S. 2000, *AJ*, 120, 3311
- Petit, J.-M., Morbidelli, A., & Valsecchi, G. B. 1999, *Icarus*, 141, 367
- Pfalzner, S. 2003, *ApJ*, 592, 986
- Quillen, A. C. & Thorndike, S. 2002, *ApJ*, 578, L149
- Quillen, A. C., Varniere, P., Minchev I., & Frank, A. 2004, *AJ*, in press
- Stern, S. A., & Colwell, J. E. 1997, *AJ*, 114, 841
- Terlevich, E. 1987, *MNRAS*, 224, 193
- Theuns, T. 1992, *A&A*, 259, 503
- Thommes, E. W., Duncan, M. J., & Levison, H. F. 2002, *AJ*, 123, 2862
- Trujillo, C. A., & Brown, M. E. 2002, *ApJ*, 566, L125
- Ward, W. R. 2003, *ApJ*, 584, L39
- Wilner, D.J., Holman, M.J., Kuchner, M. J., & Ho, P.T.P. 2002, *ApJ*, 569, L115




FIG. 1.— Orbital angles for parabolic and hyperbolic encounters, based on the similar figure by Kobayashi & Ida (2001).




FIG. 2.— Planetsesimal distribution following a retrograde hyperbolic encounter with a star. The star has $M_* = 0.2M_\odot$, orbit pericenter at $q = 40\text{AU}$, inclination $i_* = 170^\circ$, argument of perihelion $\omega_* = 20^\circ$ and eccentricity $e_* = 1.4$. The lower left shows the eccentricity distribution as a function of semi-major axis. The upper left shows the inclination distribution as a function of semi-major axis. The right hand side shows histograms for planetesimals with semi-major axes between 35 and 50 AU. This low inclination encounter scatters a significant fraction of the planetesimals, while retaining a low inclination and low eccentricity analog to the classical Kuiper Belt population.




FIG. 3.— Known Trans-Neptunian objects from the minor planet center database as of Jan 7, 2004. To ensure accurate orbits, only objects that have been observed over more than one opposition have been plotted. The known KBO distribution is displayed identically to the simulation shown in Figure 2. The known KBO distribution of orbital elements has high eccentricity and inclination tails similar to those seen in the simulation presented in Fig. 2. The known KBO distribution is affected by observational selection effects and has had 4 billion years of dynamical evolution since formation. High eccentricity objects that are not in mean motion resonances with Neptune are likely to have been removed from the Kuiper belt.




FIG. 4.— Similar to Figure 2 except the encounter is prograde. This simulation corresponds to an encounter with $M_* = 0.1M_\odot$, $e_* = 1.5$, $i_* = 20^\circ$, $\omega_* = 20^\circ$, and $q_0 = 60\text{AU}$. Prograde encounters can also cause a dichotomy in the scattered population similar to what is seen in the Kuiper Belt. However, we have failed to identify a region in parameter space that retains low eccentricity objects as well as low inclination ones, and still scatters significant fraction of the belt.




FIG. 5.— Similar to Figure 2 except the planetesimal disk prior to the encounter was truncated at 45 AU prior to the encounter. This simulation corresponds to an encounter with $M_* = 0.25M_\odot$, $e_* = 1.3$, $i_* = 170^\circ$, $\omega_* = 20^\circ$, and $q_0 = 40\text{AU}$. Following the encounter, objects initially near the edge of the disk can be scattered to higher eccentricity, and semi-major axis orbits. Past 45 AU, both the mean eccentricity and inclination increase with semi-major axis. A previously truncated disk cannot explain the eccentricity increase with semimajor axis seen in the Kuiper belt because there is no increase in mean inclination past 45 AU in the Kuiper belt.